

Multiquark states as di-hadronic molecules

Ajay Kumar Rai^{a,b}, J N Pandya^b and P C Vinodkummar^a

^aDepartment of Physics, Sardar Patel University, Vallabh Vidyanagar,
Gujarat-388 120, INDIA.

^bApplied Physics Dept., Faculty of Technology and Engineering,
M S University of Baroda, Vadodara, Gujarat-390 001, INDIA.

Multiquark systems such as tetraquarks, pentaquarks and hexaquarks states are studied as di-hadronic molecules in a nonrelativistic model. The masses of several di-hadronic states using a molecular interaction provided by asymptotic expression of the confined gluon exchange potential are computed. The exotic states such as f_0 (0.982), f_2 (1.565), f_2 (1.950), $X(3.87)$, D_{sJ} (2.317, 2.460, 2.632), $\psi(4.040)$ etc are identified as the low lying di-mesonic states, and Θ^+ (1.54) as a $K - N$ molecular state.

1. Introduction

The exotic states include those which do not fit in the $q\bar{q}$ and qqq states. So they include the tetra quark states, penta quark states, hexa quark states, hybrids such as $q\bar{q}g$ and gg or ggg glueball states. The multi-quark system has previously been studied in the framework of Bag model [1] and nonrelativistic potential models [2]. But there exist little succuss towards understanding the tetra-quark and penta-quark states as di-hadron molecules due to nonperturbative nature of QCD at the hadronic scale. It is widely believed that the QCD sum rule and the instanton based models are capable nonperturbative methods to extract the properties of hadrons [3,4] but there exist little works on multiquark states, especially their systematic calculations [5]. Recently, several new states $X(3.872)$, D_{sJ} (2.317, 2.460, 2.632), $Z(3.931)$, $Y(3.943)$, $Y(4.260)$ etc were observed in the experimental investigations of charmed mesons. These experimental observations have generated renewed interest in their theoretical interpretation [6,7,8]. Following the molecular interpretation of these states [9] we study these systems of multiquark states as di-hadronic molecules. For the binding energy of the di-hadronic state, we consider a large r ($r \rightarrow \infty$) limit of the confined gluon propagator that appear in the COGEP [10,11]. This residual confined gluon interaction conceptually is similar to the Van-der Waal interaction of the atomic molecules.

2. Masses of Di-hadrons as molecular states

The low-lying di-hadronic molecular system consisting of either di-meson tetra quark states, meson-baryon pentaquark states and baryon-baryon hexa quark states are studied

non-relativistically. Here the di-hadronic Hamiltonian is assumed as,

$$H = M + \frac{P^2}{2\mu} + V(R_{12}) + V_{SD}(S_1 S_2) \quad (1)$$

where $M = m_{h_1} + m_{h_2}$, m_{h_1} and m_{h_2} are masses of the hadrons, μ is the reduced mass, P is the relative momentum of the two hadrons and $V(R_{12})$ is the residual (molecular) interaction potential between the two hadrons. We express it as the asymptotic expression ($r \rightarrow \infty$) of the confined one gluon exchange interaction (COGEP) given by [10,11]

$$V(R_{12}) = \frac{-k_{mol}}{R_{12}} e^{-C^2 R_{12}^2/2} \quad (2)$$

where k_{mol} is the residual strength of the strong interaction coupling and C is the effective colour screening parameter of the confined gluons. Using a trial wave function given by

$$\psi(R_{12}) = \left(4 \frac{\Omega^{3/2}}{\sqrt{\pi}}\right)^{1/2} e^{-\Omega R_{12}^2/2} \quad (3)$$

the ground state energy is obtained by minimizing the expectation value of H as

$$E(\Omega) = \langle \psi(R_{12}) | H | \psi(R_{12}) \rangle \quad \text{and} \quad \frac{dE(\Omega)}{d\Omega} = 0 \quad (4)$$

With the resultant variational parameter Ω satisfying Eqn(4) the mass of the low lying di-hadronic state is obtained as

$$E(\Omega) = M + \frac{3\Omega}{4\mu} - \frac{4k_{mol}\Omega^{3/2}}{c^2 + 2\Omega} + \frac{8}{9} \frac{\alpha_s}{m_{h_1} m_{h_2}} \vec{S}_1 \cdot \vec{S}_2 |\psi(0)|^2 \quad (5)$$

Where, we have added the spin-hyperfine contribution separately. The binding energy of the di-mesons is obtained by subtracting the rest mass energies of the respective constituent hadrons as $BE = |m_{h_1} + m_{h_2} - E|$.

We have computed the masses and binding energies of the di-hadronic systems by choosing the residual interaction strength k_{mol} as the strong interaction strength, α_s at the respective hadronic scale. Accordingly for light flavour hadrons (u, d and s sector), $k_{mol}=0.7$ and for the charmed sector $k_{mol}=0.45$ are considered. The gluon confinement parameter $C=0.1$ GeV for combinations of u, d and s quarks and $C=0.250$ GeV for c quark combinations and the experimental masses of the constituting hadrons [12] are employed in the present computations.

3. Conclusion and Discussion:

We have computed the low-lying masses of the multi-quark states by considering them as di-hadronic molecules. The inter-hadronic interaction has been taken as that due to the asymptotic expression of the confined one gluon exchange interaction among the quarks [11]. While the hadronic masses have been taken from the experimental results, the Van-der-Waal like of di-hadronic binding energies have been computed using the variational approach with the inter-hadronic interaction strength, $k_{mol} = \alpha_s$, the strong interaction running coupling constant at the respective hadronic scale.

Table 1

Low-lying masses of Multiquarks as di-hadronic molecule

Systems $h_1 - h_2$	J^{PC}	Ω GeV^2	ψ $GeV^{3/2}$	BE GeV	Mass GeV	Expt[12] GeV	Others[Ref] GeV
$\pi-\pi$	0^{++}	0.0032	0.0203	0.017	0.297		
$\pi-K$	0^{++}	0.0131	0.0582	0.025	0.659	$f_0(0.600)$	
$\pi-\rho$	1^{+-}	0.0155	0.0659	0.024	0.935		
$K-K$	0^{++}	0.0603	0.1828	0.004	0.992	$f_0(0.980)$	0.987[4]
$\pi-K^*$	1^{+-}	0.0161	0.0680	0.023	1.055	$h_1(1.170)$	
$\rho-K$	1^{+-}	0.0868	1.3512	0.004	1.261	$b_1(1.235)$	
$K-K^*$	1^{+-}	0.0961	0.2593	0.006	1.380	$h_1(1.380)$	
$\rho-\rho$	0^{++}	0.1380	0.3401	0.196	1.346	$f_0(1.370)$	
	1^{++}			0.105	1.437	$f_1(1.420)$	
	2^{++}			0.076	1.618	$f_2(1.565)$	
$\rho-K^*$	0^{++}	0.1577	0.3759	0.210	1.453	$a_0(1.450)$	
	1^{++}			0.114	1.549		
	2^{++}			0.078	1.741	$f_0(1.710)$	
K^*-K^*	0^{++}	0.1822	0.4189	0.228	1.556		
	1^{++}			0.125	1.659		
	2^{++}			0.081	1.865	$f_2(1.950)$	
$\pi-D$	0^{++}	0.0186	0.0757	0.022	2.027		
$\pi-D^*$	1^{+-}	0.0188	0.0762	0.022	2.169		
$K-D$	0^{++}	0.1415	0.3465	0.015	2.344	$D_{sj}^+(2.317)$	
$K-D^*$	1^{+-}	0.1455	0.3539	0.016	2.485	$D_{sj}^+(2.460)$	
$\rho-D$	1^{+-}	0.2684	0.5602	0.033	2.603		
K^*-D	1^{+-}	0.3265	0.6489	0.039	2.718		
$\rho-D^*$	0^{++}	0.2795	0.5775	0.235	2.543		
	1^{++}			0.134	2.644		
	2^{++}			0.064	2.845		
K^*-D^*	0^{++}	0.3420	0.6718	0.158	2.624	$D_{sJ}(2.632)$	
	1^{++}			0.040	2.741		
	2^{++}			0.077	2.976		
$D-D$	0^{++}	0.3568	0.6935	0.008	3.738		3.723 [13]
$D-D^*$	1^{+-}	0.3810	0.7285	0.006	3.878	X(3.870)	3.876 [8]
D^*-D^*	0^{++}	0.4081	0.7670	0.084	3.930		
	1^{++}			0.040	3.974		
	2^{++}			0.048	4.062	$\psi(4.040)$	3.968 [6]
K-N	$(1/2)^-$	0.1910	0.4340	0.106	1.540	$\Theta^+(1.540)$	1.54 ± 0.01 [14]
$K-\Sigma$	$(1/2)^-$	0.2212	0.4845	0.102	1.785		
K- Δ	$(3/2)^-$	0.2255	0.4916	0.011	1.737		
K^*-N	$(1/2)^-, (3/2)^-$	0.3663	0.7073	0.073	1.905		
N-N	$0^+, 1^+$	0.3843	0.7332	0.039	1.918		
N- Δ	$1^+, 2^+$	0.4836	0.8712	0.053	2.225		2.17 [15]
$\Delta-\Delta$	$0^+, 1^+, 2^+, 3^+$	0.6290	1.0610	0.032	2.496		2.46 [15]

Our predictions shown in Table-1 are compared and identified with some of the experimentally known exotic hadronic states. These exotic states are those whose spin parity do not match with the expected quark anti-quark structure for mesons and 3-quark structure of baryons. Accordingly, the pseudoscalar di-mesonic and vector-vector di-mesonic combinations will have the parity and charge conjugation PC as $++$ while for the combinations of pseudoscalar-vector di-meson state will have PC value $+-$, as shown in Table-2. The experimental candidates with the predicted J^{PC} values are chosen for comparison.

We found many 0^{++} di-mesonic states in the energy range 0.297 GeV to 1.865 GeV at the light flavour (u, d, s) sector. Many of these states may be identified with experimentally known exotic mesonic states. Accordingly we identify $h_1(1.170 \text{ GeV}) 1^{+-}$ and $f_0(0.980 \text{ GeV}) 0^{+-}$ states as the $\pi - K^*$ and K-K di-mesonic states. Other di-mesonic states such as $D_{sj}^+(2.317)$, $D_{sj}^+(2.460)$, $D_{sJ}(2.632 \text{ GeV})$, $X(3.870 \text{ GeV})$ and $\psi(4.040 \text{ GeV})$ are identified here as $K - D$, $K - D^*$, $K^* - D^*$, $D - D^*$ and $2D^*$ di-mesonic states respectively. Among many combinations of mesons-baryon and di-baryon molecular states which can be studied, only few low lying state are presented here in Table-1. Though the interpretation of $\Theta^+(1.54)$ is still doubted, we identify it as K-N molecular state. Many of the other predicted di-hadronic states could be experimentally identified.

REFERENCES

1. R. L. Jaffe, *Phys. Rev. D* **38** (1977) 195.
2. J. Weinstein and N. Isgur *Phys. Rev. D* **27**, (1983) 588.
3. D. Diakonov, *Progress in Particle and Nuclear Physics* **51**(2003)173.
4. J. A. Oller and E. Oset, *Nuclear Physics A* **620** (1997) 438.
5. Jun Sugiyama et. al., arXiv : hep-ph/0309271 v 1 24 Sep.(2003)
6. D. Ebert, R. N. Faustov and V. O. Galkin, *Phys. Lett. B* **634** (2006) 214.
7. Marina Nielsen, *Phys. Lett. B* **634** (2006) 35.
8. D. Jane and M. Rosina, *Few-Body Systems* **35** (2004) 175.
9. P. Colangelo, F De Fazio and R. Ferrandes, *Mod. Phys. Lett. A* **19**, (2004) 2083; hep-ph/0407137
10. Ajay Kumar Rai, J N Pandya and P C Vinodkumar, *Indian J.Phys.* **A 80(4)** (2006) 387 (and reference therein); hep-ph/0603008.
11. J N Pandya and P C Vinodkumar, *Pramana J. Phys* **57** (2001) 821. (and reference therein).
12. S. Eidelman et. al. (Particle Data Group), *Phys. Lett. B* **592** (2004) 1 and (2005).
13. L. Maiani, F. Piccinini, A. D. Polosa, V. Riquer, *Phys. Rev. D* **71** (2005) 014028.
14. T. Nakano et. al. *Phys. Rev. Lett.* **91** (2003) 012002.
15. H. Garcilazo, A. Valcarce and F. Fernandez *Phys. Rev. C* **60** (1999) 044002 and X. Q. Yuan, Z.Y. Zhang, Y. W. Yu and P. N. Shen *ibid* (1999) 045203.